Newsletter genetic | No. 1/2014

A Service by LABOKLIN - Laboratory for clinical diagnostics www.laboklin.com





Package deals for genetic testing -

available until 30.04.2014

We are pleased to present you with a variety of genetic testing combinations as a special promotional offer. We have put together breed-specific genetic test combinations - based on the needs of our customers over the past few years. The packages contain only genetic tests that are sensible for the various breeds, since the examined gene variants have been verifiably related to disease. Therefore, our packages are not a random assortment of genetic tests, but rather sensible combinations dependent on the breed. They are put together as follows:



IOA + SCA€ 76.73 A savings of 23%1)

LOA + SCA + PLL € 97.07 A savings of 35%1)



Australian Shepherd

CEA* (Optigen) + Cataract (HSF4) + MDR* € 190.41 A savings of 18%1)

Optional prcd-PRA* can be purchased in addition



Lagotto Romagnolo

Juvenile Epilepsy + Furnishing € 76,73 A savings of 34%¹⁾

Parson Russell Terrier Australian Shepherd, Bobtail, Border Collie, German Shepherd, Elo, short- and longhaired Collie, Longhaired Whippet, McNab, Shetland Sheepdog, Silken Windhound, Waeller and White Swiss Shepherd

MDR* + DM A savings of 34% ¹⁾	€ 76,73
MDR* + DM + MH	

+ Dilution* € 115,55 A savings of 46%1)

MDR* + DM + MH+ Dilution* + HUU € 132,19 A savings of 50%1)

For all breeds

MH + Dilution* € 76.73 A savings of 23%¹

Labrador Retriever

prcd-PRA* (Optigen) + EIC* + HNPK* + DNA-Profile (ISAG 2006) € 244,97 A savings of 25%1

*patented / licensed tests

1) compared to individual orders

The prices are exclusive VAT and are valid until 30.04.2014 (date of sample arrival). To order any of these packages, please use the attached lab order form. This form can also be found on our homepage, www.laboklin.com

D-97688 Bad Kissingen · Steubenstraße 4 Tel. +49 971 72020 · Fax +49 971 68546 info@laboklin.de

CH-4058 Basel · Riehenring 173 Tel. +41 61 3196060 · Fax +41 61 3196065 labor.basel@laboklin.ch

A-4040 Linz · Rosenstraße 1 Tel. +43 732 717242 · Fax +43 732 717322 labor.linz@laboklin.at





only dogs that obtain the defective gene from both the sire and dam will

Through genetic testing, it is possible to identify LOA in Parson Russell Terriers and SCA in Parson and Jack Russell Terriers. Using selective breeding, the birth of affected dogs can be avoided.

Newsletter Archive

This and all our previous newsletters can be found on our website under the rubric "genetics" as pdf files for viewing and/or printing.

Ataxia in the Jack Russell Terrier and **Parson Russell Terrier**

The hereditary ataxia in Jack Russell Terriers and Parson Russell Terriers has been extensively studied in recent years by various research groups, making it possible to investigate the genetic causes of the various forms. At present there are three forms of ataxia each of which are caused by different mutations.

"Late Onset Ataxia" (LOA)

One form is "late onset ataxia" (LOA), which can occur in Parson Russell Terriers. An occurrence of LOA in "Early Onset Ataxia" (EOA) Jack Russell Terriers has neither been confirmed nor disqualified. The first symptoms to be observed are coordination problems of the musculoskeletal system and balance difficulties. The disease worsens in the months following onset, usually so much so that, ultimately, it ends in complete immobility.

"Spinocerebellar Ataxia" (SCA)

In "spinocerebellar ataxia" (SCA) similar symptoms as in LOA manifest, but according to experts the course of disease is more severe. Characteristic to SCA are muscle twitches or stiffness as well. Affected dogs usually do not reach three years of age.

Both forms usually occur in dogs between 3 and 6 months of age and worsen in severity in the months to follow.

A third form of ataxia, which occurs already after a few weeks of age, is called "early onset ataxia" (EOA) accordingly, and manifests mainly as a lack of coordination of head movements and front paws. The genetic cause of this form is presently not known.

The above mentioned genetically related forms of ataxia are inherited autosomal recessively. Therefore,



Any questions?

E-mail to labogen@laboklin.de

Dr. rer. nat. Petra Kühnlein Head of department Molecular Biology

Tel. +49 (0) 9 71 - 72 02 0





Download submission forms: www.laboklin.com, then go to Genetic - Lab order